

Food Safety in Pregnancy

Extension Consumer Food Safety Fact Sheet- September 2013

Pregnancy is a time when a great deal of advice is given, including some that is based on science, as well as some that is not. One area where pregnant women need to seek out scientific information is food safety. This will help keep her and her unborn baby healthy and reduce her risk of foodborne illness.



Why is food safety more important for pregnant women?

Pregnant women’s bodies naturally undergo hormonal changes which also change their immune system, making their body more susceptible to foodborne illness. Further, her unborn child has an under-developed immune system, which also puts them at greater risk. All types of foodborne illnesses are a concern for everyone, particularly pregnant women and other vulnerable populations such as the very young, old, and immune-compromised people. However, there are certain harmful microorganisms that can easily transfer from a pregnant woman to her baby which causes them to be of greater concern, including the following:

- *Listeria monocytogenes*: can lead to a disease called listeriosis, which can cause miscarriage, premature delivery, serious sickness, or even death of a newborn baby.
- *Toxoplasma gondii*: a parasite found in numerous food sources, as well as dirty cat litter boxes and other areas where cat feces can be found. Toxoplasmosis can cause infant hearing loss, mental retardation, and blindness.

What food choices can I make to lower my risk of foodborne illness?

<u>Higher risk</u>	<u>Lower risk</u>	<u>Illness prevented</u>
⚠ Raw or undercooked meat or poultry	✅ Meat and poultry cooked to a safe minimum internal temperature ¹	<i>E. coli, Salmonella</i>
⚠ Raw or undercooked fish or seafood (i.e. sashimi); refrigerated smoked fish	✅ Previously cooked seafood reheated to 165°F; canned fish and seafood; seafood cooked to 145°F ²	Parasites, bacteria, such as <i>Vibrio spp.</i>
⚠ Unpasteurized (raw) milk	✅ Pasteurized milk	<i>Listeria, E. coli, Campylobacter, Salmonella</i>
⚠ Unpasteurized (raw) fruit juice or cider	✅ Fruit juice or cider labeled “pasteurized”	<i>E. coli</i>

¹ Use a food thermometer to check for the following temperatures: Beef, pork, lamb and veal steaks, chops and roasts: 145°F with a three (3) minute rest time after cooking; ground beef, pork, lamb and veal: 160°F; all poultry products, all reheated foods: 165°F

² Pregnant women and young children should also monitor their consumption of fish containing mercury. More information is available from: www.fda.gov/Food/FoodbornellnessContaminants/BuyStoreServeSafeFood/ucm110591.htm

Æ Foods containing raw/ undercooked eggs, i.e. homemade raw cookie dough, homemade eggnog, homemade ice cream	Ü At home, use pasteurized eggs/egg products in recipes calling for raw or undercooked eggs; when eating out, ask if pasteurized eggs were used.	<i>Salmonella</i>
Æ Soft boiled or “over-easy” eggs	Ü Fully cooked eggs with firm yolk and whites; Dishes containing eggs (quiche, etc.) should be cooked to 160°F	<i>Salmonella</i>
Æ Raw sprouts (alfalfa, bean, any other)	Ü Cooked sprouts	<i>E. coli, Salmonella</i>
Æ Unwashed fresh fruits and vegetables	Ü Washed fresh fruits and vegetables; cooked vegetables	<i>Listeria, Salmonella</i>
Æ Soft cheeses made from unpasteurized (raw) milk, such as Feta, Brie, Camembert, Blue-veined, Queso fresco	Ü Hard cheeses, processed cheeses, cream cheese, soft cheeses clearly labeled “made from pasteurized milk”	<i>E. Coli, Listeria</i>
Æ Hot dogs, deli meats eaten cold (not reheated)	Ü Hot dogs, deli meats reheated to 165°F immediately before consumption	<i>Listeria</i>
Æ Unpasteurized, refrigerated pates or meat spreads	Ü Canned or shelf-stable pates or meat spreads	<i>Listeria</i>

What other food safety practices should I be following?

All consumers should follow the Four Basic Steps to Food Safety: Clean, Separate, Cook, and Chill. More information is available from: www.foodsafety.gov/keep/basics/index.html

Other useful resources:

- *Food Safety for Pregnant Women*; USDA, FDA. Available from www.fsis.usda.gov
- FDA and USDA information: www.foodsafety.gov/poisoning/risk/pregnant/
- *Nutrition for women: Special considerations and recommendations before, during and after pregnancy*: <http://missourifamilies.org/features/nutritionarticles/nut135.htm>
- Your doctor or health care provider should also have good advice and information available

Prepared by Londa Nwadike, PhD, Kansas State University/ University of Missouri Extension Food Safety Specialist

Reviewed by: - Sandy Procter, PhD, RD, LD, State Nutrition Specialist, K-State Research and Extension
 - Candance Gabel, MS, RD, LD, Associate State Nutrition Specialist, University of Missouri Extension

22201 W. Innovation Dr
 Olathe, KS 66061
 tel: 913-307-7391
 email: lnwadike@ksu.edu

1600 NE Coronado Dr
 Blue Springs, MO 64014
 tel: 816-655-6258
 email: nwadikel@missouri.edu



■ Issued in furtherance of the Cooperative Extension Work Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. Director, Cooperative Extension, University of Missouri, Columbia, MO 65211
 ■ an equal opportunity/ADA institution ■ 573-882-7216 ■ extension.missouri.edu